

## REVIEW

## Open Access

# Prevalence, incidence and risk factors for anogenital warts in Sub Saharan Africa: a systematic review and meta analysis

Cecily Banura<sup>1\*</sup>, Florence M Mirembe<sup>2</sup>, Jackson Orem<sup>3</sup>, Anthony K Mbonye<sup>4</sup>, Simon Kasasa<sup>5</sup> and Edward K Mbidde<sup>6</sup>

## Abstract

**Introduction:** The quadrivalent HPV vaccine is highly effective in primary prevention of anogenital warts (AGWs). However, there is lack of systematic review in the literature of the epidemiology of AGWs in Sub Saharan Africa (SSA).

**Objective:** To review the prevalence, incidence and risk factors for AGWs in SSA prior to the introduction of HPV vaccination programs.

**Methods:** PubMed/MEDLINE, Africa Index Medicus and HINARI websites were searched for peer reviewed English language published medical literature on AGWs from January 1, 1984 to June 30, 2012. Relevant additional references cited in published papers were also evaluated for inclusion. For inclusion, the article had to meet the following criteria (1) original studies with estimated prevalence and/or incidence rates among men and/or women (2) detailed description of the study population (3) clinical or self-reported diagnosis of AGWs (4) HPV genotyping of histologically confirmed AGWs. The final analysis included 40 studies. Data across different studies were synthesized using descriptive statistics for various subgroups of females and males by geographical area. A meta - analysis of relative risk was conducted for studies that had data reported by HIV status.

**Results:** The prevalence rates of clinical AGWs among sex workers and women with sexually transmitted diseases (STDs) or at high risk of sexually transmitted infection (STIs) range from 3.3% - 10.7% in East, 2.4% - 14.0% in Central and South, and 3.5% - 10.5% in West African regions. Among pregnant women, the prevalence rates range from 0.4% - 3.0% in East, 0.2% - 7.3% in Central and South and 2.9% in West African regions. Among men, the prevalence rates range from 3.5% - 4.5% in East, 4.8% - 6.0% in Central and South and 4.1% to 7.0% in West African regions. In all regions, the prevalence rates were significantly higher among HIV+ than HIV- women with an overall summary relative risk of 1.62 (95% CI: 1.43-1.82). The incidence rates range from 1.1 - 2.7 per 100 person-years among women and 1.4 per 100 person years among men. Incidence rate was higher among HIV+ (3.0 per 100 person years) and uncircumcised men (1.7 per 100 person-years) than circumcised men (1.3 per 100 person-years).

HIV positivity was a risk factor for AGWs among both men and women. Other risk factors in women include presence of abnormal cervical cytology, co-infection with HPV 52, concurrent bacteria vaginosis and genital ulceration. Among men, other risk factors include cigarette smoking and lack of circumcision.

**Conclusions:** AGWs are common among selected populations particularly HIV infected men and women. However, there is need for population-based studies that will guide policies on effective prevention, treatment and control of AGWs.

**Keywords:** Anogenital warts, Sub Saharan Africa, HIV, HPV vaccination

\* Correspondence: [cbanura@chdc.mak.ac.ug](mailto:cbanura@chdc.mak.ac.ug)

<sup>1</sup>Department of Child Health and Development Centre, Makerere University College of Health Sciences, Kampala, Uganda

Full list of author information is available at the end of the article

## Introduction

The epidemiology of AGWs in most of SSA is largely unknown since few studies have been conducted. Studies from high income countries show that the clinical burden has been increasing over the years since approximately 0.5-1.0% of adults below 50 years have AGWs [1-3].

Caused mainly by low-risk HPV type 6 and 11, AGWs affect both men and women [1]. They are highly infectious with about 65% of individuals with an infected partner developing lesions within 3 weeks [4]. The median time between infection and development of lesions is about 11–12 months among men [5,6] and about 5–6 months among women [7]. Rarely, AGWs have been associated with malignant Bushke-Lowenstein malignant tumors [8]. Their occurrence is strongly linked to HIV and weakly associated with cigarette smoking [1,4,5]. At present, the impact of highly active anti-retroviral therapy (HAART) remains unclear [5,6].

Except for Rwanda [9], most countries in SSA are yet to introduce or scale up HPV vaccination in national immunization programs. While reduction in disease burden due to HPV 16/18 may not be evident for decades, vaccination with the quadrivalent HPV vaccine should result in immediate measurable reduction in the incidence rate of AGWs. Preliminary results from the Australian national HPV vaccination program shows a significant decline in the number of cases of AGWs among the young vaccinated women and some herd-immunity effect in young unvaccinated heterosexual men [10]. This review was undertaken to assess the prevalence, incidence and risk factors for AGWs before the introduction of HPV vaccination programs in order to provide a basis for future program evaluation in SSA.

## Methods

### Identification and eligibility of relevant studies

PubMed/MEDLINE, Africa Index Medicus and HINARI websites were searched for peer reviewed English language published medical literature from January 1, 1984 up to June 30, 2012. The following Medical Subject Heading (MESH) and search terms were used alone or in combination “Sub Saharan Africa” AND (“anogenital warts” OR “venereal warts” OR “condylomata acuminata” OR “condylomata”) AND “risk factors”. Relevant additional references cited in published papers were also evaluated for inclusion. For inclusion, articles had to meet the following criteria (1) original studies with clear estimation of prevalence and/or incidence rates among men and/or women (2) detailed description of study population (3) clinical or self-reported diagnosis of AGWs (4) HPV genotyping of histologically confirmed AGWs. Studies focusing exclusively on case reports and commentaries were excluded.

### Data extraction

From each article, the following information was extracted: first author, publication journal name and year of publication, country of study population, study sample type (population- or clinic- based), study design, mean or median age with range/inter quartile range, whenever available, sample size, prevalence and/or incidence rate overall and by HIV status, whenever available, risk factors and the overall prevalence of HIV, whenever available. For studies that included populations from different countries, data was extracted separately for each country.

### Statistical analysis

Data across different studies were synthesized using descriptive statistics for different subgroups of females and males by geographical areas. A meta- analysis of relative risk was conducted for studies that had data reported by HIV status and results presented in a forest plot. In total, 40 studies (39 hospital - and 1 population-based) are included in this review.

## Results

The prevalence and incidence rates of AGWs in diverse female and male hospital-based study populations in East, Central and South, and West African regions is summarized in Tables 1, 2 and 3. Overall, there is inter- and intra-region variations in rates depending on the underlying prevalence rate of HIV-1 infection, study population and age group studied. While both young and adult populations were studied, there seems to be no trend or pattern of prevalence rates by age.

### Prevalence rates among women by geographical region

The prevalence rates among women in East, Central and South, and West African regions are summarized in Tables 1. The prevalence rates of clinical AGWs among sex workers and women with STDs or at high risk for other STIs range from 3.3% - 10.7% in East, 2.4% - 14.0% in Central and South, and 3.5% - 10.5% in West African regions. Among pregnant women, the prevalence rates range from 0.4% - 3.0% in East, 0.2% - 7.3% in Central and South and 2.9% (single study) in West African regions.

### Prevalence rates of AGWs among men by geographical region

Eight (8) studies (3 East Africa, 2 Central and South Africa, 3 West Africa) reported prevalence rates of AGWs in men (Table 2). The prevalence rate among STD clinic attendees, men who have sex with men, and men with symptoms of STDs in Central and South and West African region range from 4.8% - 12.2% and 2.8% - 4.1%,

**Table 1 Studies reporting prevalence of AGWs in women**

Author, Publication year	Country	Study design	Study population	Sample size	Mean or Median age (years, range/IQR <sup>1</sup> )	Prevalence of AGWs <sup>2</sup> n (%)	Prevalence of HIV-1 (%)	Comments
<b>East Africa</b>								
Kreiss et al., 1992 [11]	Kenya	<sup>5</sup> Cross-sectional	Sex workers	196	30.2 (HIV-1+) 31.5 (HIV-1-)	18/196 (9.2) Overall 15/145 (10.0) HIV-1+ 3/51 (6.0) HIV-1		
Fonck et al., 2000 [12]	Kenya	"EntryTbl_st" <sup>5</sup> Cross-sectional	Women attending STD <sup>3</sup> clinic	520	26 ± 6.8 (14–49)	31/520 (6.0) 5/520 (1.0) <sup>a</sup>	29.0	Prevalence of AGWs 5% (Non pregnant women) 9% (Pregnant women) 6% (One sexual partner)
Mayaud et al., 2001 [13]	Tanzania	<sup>5</sup> Cross-sectional	Pregnant women	660	23.4 ± 5.1 (15–44)	20/660 (3.0)	15.0	
Riedner et al., 2003 [14]	Tanzania	<sup>5</sup> Open cohort	Female bar workers	600	25.4	39/600 (6.5) Overall 39/408 (9.6) HIV + 0/192 (0.0) HIV -	68.0	
Namkinga et al., 2005 [15]	Tanzania	<sup>5</sup> Cross-sectional	Women presenting with complaints of genital infections	464		18/464 (3.9)	22.0	
Amone-P'Olak, 2005 [16]	Uganda	<sup>4</sup> Cross-sectional	Formally abducted teenage girls in Northern Uganda	123	16.2 ± 2.2 (12–18)	67/123 (54.5) <sup>a</sup>		
Mbizvo et al., 2005 [17]	Tanzania	<sup>5</sup> Cross-sectional	Women seeking primary health care services	382	26.7 ± 6.0	8/382 (2.1)	11.5	
Msuya et al., 2006 [18]	Tanzania	<sup>5</sup> Cross-sectional	Women seeking reproductive health care services	382	24.6 (14–43)	7/382 (2.0)	6.9	
Riedner et al., 2006 [19]	Tanzania	<sup>5</sup> Serial cross-sectional	Female bar workers	600	25.5 (16–39)	5.2-10.7	67.0	
Aboud et al., 2008 [20]	TanzaniaMalawi and Zambia	<sup>5</sup> Cross-sectional	HIV-1 positive pregnant women	2292	(15–49)	195/2292 (8.5)		Prevalence of AGWs Blantyre – 42/474 (8.9) Lilongwe – 61/748 (8.2) Dar es Salaam – 31/428 (7.2) Lusaka – 61/642 (9.5)
Banura et al., 2008a [21]	Uganda	Baseline of a prospective cohort study	Young women attending a clinic for teenagers	1275	20 (12–24)	97/1275 (7.6)	8.6	
Banura et al., 2008b [22]	Uganda	<sup>5</sup> Baseline of a prospective cohort study	Pregnant women Attending ANC <sup>5</sup>	987	19 (14–24)	61/987 (6.2)	7.3	
Urassa et al., 2008 [23]	Tanzania	<sup>5</sup> Cross-sectional	Youth attending an STI <sup>4</sup> clinic	214	20.2 (Females) (13–24) 21.5 (Males) (11–24)	7/214 (3.3)	15.3	HIV –1 prevalence in Males – 7.5%
Grijzen et al., 2008 [24]	Kenya	<sup>5</sup> Baseline of a prospective cohort study	Women at risk for HIV-infection	361	27 (23–32)	8/361 (2.4)	32.0	
Msuya et al., 2009 [18]	Tanzania	<sup>5</sup> Cross-sectional	Pregnant women	2655	24.6 (14–43)	11/2555 (0.4) Overall 2/184 (1.1) HIV + 9/2470 (0.4) HIV -	6.9	

**Table 1 Studies reporting prevalence of AGWs in women (Continued)**

Mapingure, et al., 2000 [25]	Tanzania	<sup>5</sup> Cross-sectional	Pregnant women	2654	24.6	34/2654 (1.3) 48/2654 (1.8) <sup>b</sup>	6.9	
<b>Central and South Africa</b>								
Latif et al., 1984 [26]	Zimbabwe	<sup>5</sup> Cross-sectional	Pregnant women attending STD clinic	175	22.3	23/175 (13.7)		
Mason et al., 1990 [27]	Zimbabwe	<sup>5</sup> Cross-sectional	Women attending STD clinic	100	(15–45)	14/100 (14.0) 1/59 (1.7) <sup>a</sup>		
Kristensen 1990 [28]	Malawi	<sup>5</sup> Cross sectional	Adult women with symptoms of STIs	16,218	26.8 ± 7.5	32/16,218 (0.2)	62.4	
Nzila et al., 1991 [29]	Democratic Republic of Congo	<sup>5</sup> Cross-sectional	Female sex workers	1233		30/1233 (2.4) Overall 21/431 (5.0) HIV + 8/802 (1.0) HIV-	35.0	
Le Bacq et al., 1993 [30]	Zimbabwe	<sup>5</sup> Cross-sectional	New STD clinic attendees	146		19/146 (13.0)	69.0	
Maher et al., 1995 [31]	Malawi	<sup>5</sup> Cross-sectional	Female patients in general medical care	61	31 (16–65)	6/61 (9.8)		
Taha et al., 1998 [32]	Malawi	<sup>5</sup> Serial cross-sectional surveys	Pregnant women	1990 – 6603 HIV + 1502 HIV- 5101 1993 – 2161 HIV + 694 HIV- 1457 1995 – 808 HIV + 808 HIV- 701		1990 1993 1995 Overall 4.8 3.1 2.5 HIV + 8.3 6.3 2.7 HIV- 2.2 1.7 1.0	23.0 (1990) 30.1 (1993) 32.6 (1995)	
Klaskala et al., 2005 [33]	Zambia	<sup>5</sup> Cross-sectional	Pregnant women	3160	25 ± 5.3 (14–43)	203/3160 (6.2)		
Mbizvo et al., 2005 [17]	Zimbabwe	<sup>5</sup> Cross –sectional	Women recruited from primary health care centers	386	26.5 ± 6.8	13/386 (3.4)	29.3	
Kurewa et al., 2010 [34]	Zimbabwe	<sup>5</sup> Cross-sectional	Pregnant women	691	24.2 ± 5.1	48/691 (7.0) 50 /691 (7.3) <sup>a</sup>	25.6	
Mapingure et al., 2010 [26]	Zimbabwe	<sup>5</sup> Cross-sectional	Pregnant women	691	24.2 ± 5.1	50/691 (7.3) 33/691 (4.8) <sup>b</sup>	25.6	
Menendez et al., 2010 [35]	Mozambique	<sup>5</sup> Cross- sectional	Women attending ANC and FP <sup>5</sup> clinics and community	262	(14–61)	13/262 (5.0)	12.0	Prevalence of HIV-1 21.0% among FP clinic attendees
<b>West Africa</b>								
Oni et al., 1994 [36]	Nigeria	<sup>5</sup> Cross-sectional	STD clinic attendees	116		12/116 (10.5)		
Ghys et al., 1995 [37]	Ivory Cost	<sup>5</sup> Cross sectional	Female sexual workers	1209		105/1209 (8.7) Overall 79/567 (14.0) HIV + 26/642 (4.0) HIV -	80.0	
Meda et al., 1997 [38]	Burkina Faso	<sup>5</sup> Cross –sectional	Women attending ANC	645	25.3 ± 2.9 (15–41)	19/645 (2.9)		
Okesola et al., 2000 [39]	Nigeria	<sup>5</sup> Cross-sectional	Patients attending an STD clinic	861	(17–74)	68/861 (8.0)		

**Table 1 Studies reporting prevalence of AGWs in women (Continued)**

Bakare et al., 2002 [40]	Nigeria	<sup>5</sup> Cross-sectional	CSWs <sup>7</sup> and women without symptoms of STIs			6.5 36.4 <sup>c</sup>	34.3	
Domfeh et al., 2008 [41]	Ghana	<sup>5</sup> Cross-sectional	Women attending gynecological clinic	75	33.3 ± 9.2 (19–57)	4/75 (5.3) <sup>a</sup>		
Sagay et al., 2009 [42]	Nigeria	<sup>5</sup> Cross-sectional	Female sex workers	374	27.8 ± 6.7 (16–63)	17/374 (4.5)		Prevalence of AGWs 5/81 (6.1%) Lemon users 12/293 (4.1%) Non Lemon users
Jombo et al., 2009 [43]	Nigeria	<sup>5</sup> Cross-sectional	Patients with genital ulcer disease	699		369/699 (52.8) Overall 285/506 (56.4) HIV + 84/193 (43.6) HIV –		Prevalence <i>Males</i> : 13/329 (2.6%) <i>Females</i> : 8/177 (1.6%)
Low et al., 2011 [44]	Burkina Faso	<sup>5</sup> Baseline of Prospective cohort	CSWs and other women with high-risk sexual behaviors	765	28 (15–54)	27/765 (3.5) Overall 19/273 (7.0) HIV –1 + 8/492 (1.6) HIV –	34.9 HIV-1 0.7 HIV-1 &2	No prevalent AGWs among women on HAART

<sup>a</sup> self-reported prevalence; <sup>b</sup> self-reported prevalence for the last 12 months; <sup>c</sup> self-reported prevalence among commercial sexual workers; <sup>1</sup>Inter quartile range; <sup>2</sup>Anogenital warts; <sup>3</sup>Sexually transmitted disease;

<sup>4</sup>Sexually transmitted infection; <sup>5</sup>Antenatal care; <sup>6</sup>Family planning; <sup>7</sup>Commercial sexual workers; <sup>8</sup> hospital-based study; <sup>9</sup> Teenagers in an institution.

**Table 2 Studies reporting AGWs in men**

Author, year	Country	Study design	Study population	Sample size	Mean or Median age (years, range/IQR <sup>1</sup> )	Prevalence of AGWs <sup>2</sup> (%)	Prevalence of HIV-1%	Comments
<b>East Africa</b>								
Grijzen et al., 2008 [24]	Kenya	<sup>§</sup> Baseline of a prospective cohort study	Men at high-risk for HIV infection	536	27 (24–33)	9/500 (1.8)	21.0	
Smith et al., 2010 [45]	Kenya	<sup>§</sup> Baseline of RCT <sup>3</sup> on male circumcision	HIV negative sexually active men	2168	20 (19–28)	12/2168 (0.6) Overall 10/1089 (0.9) HIV + 2/1079 ( 0.2) HIV-		
Tobian et al., 2012 [46]	Uganda	<sup>†</sup> Cross-sectional	Heterosexual men	1399	15–49	23/1399 (1.6) <sup>a</sup> Overall 16/421 (3.8) <sup>a</sup> HIV + 7/978 (0.7) <sup>a</sup> HIV –		
<b>Central and South Africa</b>								
Le Bacq et al., 1993 [31]	Zimbabwe	<sup>§</sup> Cross-sectional	New STD clinic attendees	319		39/319 (12.2)	61.0	
Maher et al. 1995 [32]	Malawi	<sup>§</sup> Cross-sectional	In-patient male patients in general medical care	62	39 (20–90)	3/62 (4.8)		
Machekano et al., 2000 [47]	Zimbabwe	<sup>§</sup> Baseline of prospective cohort study	Male factory workers who reported symptoms of STDs	374		22/374 (6.0)	20	
Müller et al., 2010 [48]	South Africa	<sup>§</sup> Cross-sectional	Heterosexual men attending sexual health services	214	29.8 ± 7.5	108/214 (50.5)	49.5	
<b>West Africa</b>								
Okesola et al., 2000 [40]	Nigeria	<sup>§</sup> Cross-sectional	STD <sup>2</sup> clinic attendees	1,373	17–74	4.1		
Wade et al., 2005 [49]	Senegal	<sup>§</sup> Cross sectional	Men who have sex with men	463	18–52	13/463 (2.8)	18.1	21.5% Overall 0.5% HIV-2 2.9% HIV-1 & HIV-

<sup>a</sup> Self-reported prevalence.

<sup>1</sup> Inter Quartile Range.

<sup>2</sup> Commercial sexual workers.

<sup>3</sup> Randomized Controlled Trial.

<sup>§</sup> hospital-based study.

<sup>†</sup> Population-based study.

**Table 3 Studies reporting incidence rates of AGWs in men and women**

Author, year	Country	Study design	Study population and site	Sample size	Mean or median age (years, range)	Incidence rate/100 person-years of AGWs <sup>2</sup>	HIV -1 prevalence%	Comments
<b>East Africa</b>								
Lavreys et al., 1999 [50]	Kenya	Prospective cohort	HIV negative truck drivers in Mombasa	746	26 <sup>a</sup> (17–58) 29 <sup>b</sup> (16–62)	1.4 overall 1.7 Uncircumcised 1.3 Circumcised		Annual incidence of HIV-1 – 3.0%
<b>West Africa</b>								
Ozumba et al., 1991 [51]	Nigeria	Retrospective cohort (1976–85)	Female STD <sup>1</sup> clinic attendees	45	21 ( 5–36)	2.7 (range:1.6 – 3.6)		AGWs incidence highest among teenagers and students
Low et al., 2011 [44]	Burkina Faso	Prospective cohort	Female sex workers and other women at high risk	765	28 (15–54)	1.1 HIV -	34.9	HIV- 1 & HIV-2 prevalence 0.7%

<sup>a</sup> uncircumcised men.

<sup>b</sup> circumcised men.

<sup>1</sup>Sexually Transmitted Diseases.

<sup>2</sup>Anogenital warts.

respectively. The rates among men in the East African region range from 0.6 -1.8 percent.


#### Prevalence rates by HIV status

The prevalence rates of AGWs were significantly higher among HIV+ than HIV- women in all regions with an overall summary relative risk of 1.62 (95% CI: 1.43–1.82) (Figure 1). Similarly among men, clinical and self -reported

prevalence rates were higher among HIV+ than HIV- men (Table 2).

#### Incidence rates of AGWs among men and women

Only 3 studies (2 among females and 1 among males) reported incidence rates of AGWs (Table 3). The incidence rates range from 1.1 – 2.7 per 100 person-years among women and 1.4 per 100 person years among





men. The incidence rate was higher among uncircumcised (1.7 per 100 person-years) than circumcised men (1.3 per 100 person-years) [44].

#### HPV 6 and/or 11 in AGWs

Only 3 studies reported the prevalence of HPV 6 and/or 11 in biopsy specimens or swabs taken from AGWs. HPV 11 was detected in 100% vulval-vaginal wart specimens obtained from 9 prepubescent South African girls [52]. HPV 6 and/or 11 was detected in 96.3% of 108 genital swabs taken from heterosexual men with AGWs attending sexual health clinics in South Africa [48]. Among 74 specimens taken from penile warts of HIV+ men in South Africa, HPV 6 was detected in 42.5%, HPV 11 in 32.9% and HPV 6/11 in 68.5% [53].

#### Risk factors for AGWs

Only 2 studies (one among women and another among men) reported on risk factors for AGWs. Among women, the risk of prevalent AGWs was 5 times higher among HIV-1+ than HIV-1- women and 3 times higher among women who smoked cigarettes than those who did not. Among HIV-1+ women with low CD4+ count ( $\leq 200$  cells/ $\mu$ L), the risk of incident AGWs was elevated 20 fold, and 6fold for women with CD4+ count  $>200$  cells/ $\mu$ L. Other risk factors for incident AGWs in women include detection of HPV 6, concurrent bacterial vaginoses, genital ulceration, presence of abnormal cervical cytology and the detection of cervical HPV 52 [44]. Lack of circumcision and HIV infection were risk factors for AGWs in men [45].

#### Discussion

To the best of our knowledge this is the first systematic review of the epidemiology of AGWs in SSA. The literature suggests that AGWs are prevalent among both men and women populations seeking care in their respective health care systems. The fewer studies among men is not surprising given that women generally have more frequent contact with the health care system than men. Although there is no marked difference between regions, absence of a standardized protocol for diagnosis might have contributed to the observed variations across studies within the same region. Overall, the prevalence rates were higher than those reported from retrospective administrative databases or medical chart reviews in high income countries possibly because of underlying HIV infection in several studies [54].

Consistent with published studies, the risk for AGWs was higher for HIV+ than HIV- men and women [55]. HIV+ women had almost 2 fold risk for HPV infection than HIV-women. While some AGWs may have been a result of new infections, recrudescence of existing HPV infection has been reported among sexually inactive HIV+ women [56].

Impaired CD4+ T-lymphocyte response and other forms of immune dysfunction may be responsible for altering the natural history of HPV infection among HIV infected individuals [57]. The use of highly active anti-retroviral therapy has been shown to reduce the risk of opportunistic malignancies such as Kaposi sarcoma among HIV+ individuals [58], however, their impact on AGWs remains unclear [55,57,59]. On the other hand consistent use of male condoms appears to reduce the risk by 60-70% [60].

Consistent with other studies, HPV 6 and 11 alone or in combination were detected in the few studies that examined HPV genotypes in AGWs specimens albeit small sample sizes. However, the contribution of HPV 11 to the development of AGWs remains unclear [4,7]. The concurrent detection of HPV 52 with HPV 6 was not surprising as co-infection with high risk HPV types has been reported in 20-50% of AGWs [61,62].

In the absence of a clinical test to establish sub clinical HPV 6 and 11 infections, identification of risk factors for acquisition of AGWs independent of other STDs is complex. Consistent with other studies, low CD4+ cell count ( $\leq 200$  cells/ $\mu$ L) and abnormal cervical and anal dysplasias are risk factors for AGWs in HIV+ women and men, respectively [63,64]. Other risk factors for AGWs in women identified in this review included co-infection with HPV 52, and concurrent bacteria vaginoses [65]. In men, anal HPV infection and related dysplasias [39] and lack of circumcision [45] were additional risk factors.

Although AGWs are not life threatening, they cause significant psychological distress and are refractory to conventional therapies, hence the need for prevention [4,66]. The quadrivalent HPV vaccine, correct and consistent condom use and limiting the number of sexual partners are some of the prevention options available to reduce the risk of contracting AGWs.

It is important to note that there are limitations to this study. This review focused only on peer reviewed English language articles published from a few SSA countries, which limits generalization of the findings. Secondly, most studies were conducted in hospital-based study populations, which would favor higher rates than in the general population. Thirdly, the rates should be interpreted with caution because of the differences in study populations and age group studied. While some studies included all adults [31,39], others focused on narrow age ranges of specific populations like young people and pregnant women [23,25] that could have resulted in the observed high rates. Nevertheless, the review provides vital baseline data against which the impact of HPV vaccination could be evaluated in future.

#### Conclusions

AGWs are common among selected populations particularly HIV+ men and women. However, there is need



for population-based studies on AGWs that will guide policies on effective prevention, treatment and control services.

# Competing interests

The authors declare that they have no competing interests.

# Authors' contributions

CB conceived the study, searched the literature, drafted the manuscript and produced the final tables FMM, JO, AKB, SK, EKM made substantial contributions to the manuscript and contributed to data interpretation. All authors read and approved the final manuscript.

# Author details

<sup>1</sup>Department of Child Health and Development Centre, Makerere University College of Health Sciences, Kampala, Uganda. <sup>2</sup>Department of Obstetrics and Gynecology, Makerere University College of Health Sciences, Kampala, Uganda. <sup>3</sup>Uganda Cancer Institute, P.O. Box 3935, Kampala, Uganda. <sup>4</sup>Department of Community Services, Ministry of Health, P.O. Box 7272, Kampala, Uganda. <sup>5</sup>School of Public Health, Makerere University College of Health Sciences, Kampala, Uganda. <sup>6</sup>Uganda Virus Research Institute, P.O. Box 49, Entebbe, Uganda.

Received: 18 August 2012 Accepted: 1 July 2013

Published: 10 July 2013

# References

- Lacey C: Therapy for human papillomavirus-related disease. *J Clin Virol* 2005, **32**(Suppl 1):S82-S90.
- Kjaer SK, Tran T, Sparen P, Tryggvadottir L, Munk C, Dasbach E, Liaw KL, Nygård J, Nygård M: The burden of genital warts: a study of nearly 70,000 women from the general female population in the 4 Nordic countries. *J Infect Dis* 2007, **196**:1447-1454.
- Kliwer EV, Demers AA, Elliott L, Lotocki R, Butler JR, Brisson M: Twenty-year trends in the incidence and prevalence of diagnosed anogenital warts in Canada. *Sex Transm Dis* 2009, **36**(6):380-386.
- Lacey C, Lowndes C, Shah KV: Chapter 4: Burden and management of non-cancerous HPV-related condition: HPV 6/11. *Vaccine* 2006, **24**(S3):S3/S3-41.
- Arimi Y, Winer RL, Feng QA, Hughes JP, Lee SK, Stern ME, O'Reilly SF, Koutsky LA: Development of genital warts after incident detection of human papillomavirus infection in young men. *J Infect Dis* 2010, **202**:1181-1184.
- Anic GM, Lee GH, Stockwell H, Rollison DE, Wu Y, Papenfuss MR, Villa LL, Lazcano-Ponce E, Gage C, Silva RJ, Baggio ML, Quiterio M, Salmeron J, Abrahamsen M, Giuliano AR: Incidence and human papillomavirus (HPV) type distribution of genital warts in a multinational cohort of men: The HPV in men study. *J Infect Dis* 2011, **204**:1886-1892.
- Garland SM, Stenben M, Sings HL, James M, Lu S, Railkar R, Barr E, Haupt RM, Joura EA: Natural history of genital warts: analysis of the placebo arm of 2 randomized phase III trials of a quadrivalent human papillomavirus (types 6, 11, 16, and 18) vaccine. *J Infect Dis* 2009, **199**:805-814.
- Giuliano AR, Tortolero-Luna G, Ferrer E, Burchell AN, De Sanjose S, Kjaer SK, Munoz N, Schiffman M, Bosch FX: Epidemiology of human papillomavirus infection in men, cancers other than cervical and benign conditions. *Vaccine* 2008, **26**(Suppl 10):K17-K28.
- The Lancet: Financing HPV vaccination in developing countries. *Lancet* 2011, **377**(9777):1544.
- Donovan B, Franklin N, Guy R, Grulich AE, Regan DG, Ali H, Wand H, Fairley CK: Quadrivalent human papillomavirus vaccination and trends in genital warts in Australia: analysis of national sentinel surveillance data. *Lancet Infect Dis* 2011, **11**:39-44.
- Kreiss J, Kiviat N, Plummer F, Roberts P, Waiyaki P, Ngugi E, Holmes KK: Human Immunodeficiency Virus, Human papillomavirus, and cervical intraepithelial neoplasia in Nairobi prostitutes. *Sex Transm Infect* 1992, **19**(1):54-59.
- Fonck K, Kidula N, Kirui P, Ndinya-Achola J, Bwayo J, Claeys P, Temmerman M: Pattern of sexually transmitted diseases and risk factors among women attending an STD referral clinic in Nairobi, Kenya. *Sex Transm Dis* 2000, **7**:417-423.
- Mayaud P, Weiss H, Lacey C, Uledi E, Kopwe LT, Ka-Gina G, Grosskurth H, Hayes RJ, Mabey DC, Lacey CJ: The interrelation of HIV, cervical human papillomavirus, and neoplasia among antenatal clinic attenders in Tanzania. *Sex Transm Infect* 2001, **77**(4):248-254.
- Riedner G, Hoffmann O, Nichombe F, Lyamuya EM, Mmbando D, Maboko L, Hay P, Todd J, Hayes R, Hoelscher M, Grosskurth H: Baseline survey of sexually transmitted infections in a cohort of female bar workers in Mbeya region, Tanzania. *Sex Transm Infect* 2003, **79**:382-387.
- Namkinga L, Matee M, Kivaisi A, Moshiri C: Prevalence and risk factors for vaginal candidiasis among women seeking primary care for genital infections in Dar es Salaam, Tanzania. *East Afr Med J* 2005, **82**(3):138-143.
- Amone-P'Olak: Psychological impact of wae and sexual abuse on adolescent girls in northern Uganda. *Intervention* 2005, **3**(1):33-45.
- Mbizvo E, Msuya S, Hussain A, Chirenje M, Mbizvo M, Sam N, Stray-Pedersen B: HIV and sexually transmitted infections among women presenting at urban primary health care clinics in two cities of sub-Saharan Africa. *Afr J Reprod Health* 2005, **9**(1):88-98.
- Msuya S, Mbizvo E, Stray-Pedersen B, Sundby J, Sam N, Hussain A: Risk assessment at the primary health care level in Moshi, Tanzania: limits in predicting sexually transmitted infections among women. *Cent Afr J Med* 2006, **52**(9-12):97-104.
- Riedner G, Rusizoka M, Mmbando D, Maboko L, Grosskurth H, Todd J, Hayes R, Hoelscher M: Decline in sexually transmitted infection prevalence and HIV incidence in female barworkers attending prevention and care services in Mbeya Region. *Tanzania. AIDS* 2006, **20**(4):609-615.
- About S, Msamanga G, Read JS, Mwatha A, Chen YQ, Potter D, Potter D, Valentine M, Sharma U, Hoffmann I, Taha TE, Goldenberg RL, Fawzi WW: Genital tract infections among HIV-infected pregnant women in Malawi, Tanzania and Zambia. *Int J STD AIDS* 2008, **19**(12):824-832.
- Banura C, Franceschi S, van Doorn L, Arslan A, Kleter B, Wabwire-Mangen F, Mbide EK, Quint W, Weiderpass E: Infection with human papillomavirus and HIV among young women in Kampala, Uganda. *J Infect Dis* 2008, **197**(4):555-562.
- Banura C, Franceschi S, van Doorn L, Arslan A, Kleter B, Wabwire-Mangen F, Mbide EK, Quint W, Weiderpass E: Prevalence, incidence and clearance of human papillomavirus infection among young primiparous pregnant women in Kampala, Uganda. *Int J Cancer* 2008, **123**(9):2180-2187.
- Urassa W, Moshiri C, Chalamilla G, Mhalu F, Sandstrom E: Risky sexual practices among youth attending a sexually transmitted infection clinic in Dar es Salaam, Tanzania. *BMC Infect Dis* 2008, **8**:159.
- Grijns ML, Graham S, Mwangome M, Githua P, Mutimba S, Wamuyu L, Okuku H, Price MA, McClelland RS, Smith AD, Sanders EJ: Screening for genital and anorectal sexually transmitted infections in HIV prevention trials in Africa. *Sex Transm Infect* 2008, **84**:364-370.
- Mapingure MP, Msuya S, Kurewa NE, Mujoma MW, Sam N, Chirenje MZ, Rusakaniko S, Saugstad LF, de Vlas SJ, Stray-Pedersen B: Sexual behavior does not reflect HIV-1 prevalence differences: a comparison study of Zimbabwe and Tanzania. *J International AIDS Society* 2010, **13**(45).
- Latif A, Bvumbe J, Muongerwa J, Paraiwa E, Chikosi W: Sexually transmitted diseases in pregnant women in Harare, Zimbabwe. *Afr J Sex Transm Dis* 1984, **1**(1):21-23.
- Mason P, Gwanzura L, Latif A, Marowa E: Genital infections in women attending a genito-urinary clinic in Harare, Zimbabwe. *Genitourin Med* 1990, **66**:178-181.
- Kristensen J: The prevalence of symptomatic sexually transmitted diseases and human immunodeficiency virus infection in outpatients in Lilongwe, Malawi. *Genitourin Med* 1990, **66**(4):244-246.
- Nzila N, Laga M, Thiam MA, Mayimona K, Eddi B, Van Dyck E, Behets F, Hassig S, Nelson A, Mokwa K, et al: HIV and other sexually transmitted diseases among female prostitutes in Kinshasa. *AIDS* 1991, **5**:715-721.
- Le Bacq F, Mason P, Gwanzura L, Robertson V, Latif A: HIV and other sexually transmitted diseases at a rural hospital in Zimbabwe. *Genitourin Med* 1993, **69**:352-356.
- Maier D, Hoffman I: Prevalence of genital infections in medical inpatients in Blantyre, Malawi. *J Infect* 1995, **31**:77-78.
- Taha T, Dallabetta GA, Hoover DR, Chipangwi JD, Mtshayalye LA, Liomba GN, Kumwenda NI, Miotti PG: Trends of HIV-1 and sexually transmitted diseases among pregnant and post-partum women in urban Malawi. *AIDS* 1998, **12**:197-203.
- Skalka W, Brayfield BP, Kankasa C, Bhat G, West JT, Mitchell CD, Wood C: Epidemiological Characteristics of Human Herpesvirus-8 Infection in a

- Large Population of Antenatal Women in Zambia. *J Med Virol* 2005, **75**:93–100.
34. Kurewa N, Mapingure M, Munjoma M, Chirenje M, Rusakaniko S, Stray-Pedersen B: **The burden and risk factors for sexually transmitted infections and reproductive tract infections among pregnant women in Zimbabwe.** *BMC Infect Dis* 2010, **10**:127.
  35. Menendez C, Castellsague X, Renom M, Sacarlal J, Quinto L, Lloveras B, Klaustermeier J, Kornegay JR, Sigauque B, Bosch FX, Alonso PL: **Prevalence and risk factors of sexually transmitted infections and cervical neoplasia in women from a rural area of Southern Mozambique.** *Infect Dis Obstet Gynecol* 2010, doi:10.1155/2010/609315. Published online 2010 July 11.
  36. Oni A, Adu F, Ekweozor C: **Isolation of herpes simplex virus from sexually transmitted disease patients in Ibadan, Nigeria.** *Sex Transm Dis* 1994, **21**(4):187–190.
  37. Ghys PD, Diallo OM, Ettiègne-Traoré V, Yeboue KM, Gnaore E, Lorougnon F, Kal K, Van Dyck E, Brattegaard K, Hoyi YM, Whitaker JP, De Cock KM, Greenberg AE, Piot P, Laga M: **Genital ulcers associated with human immunodeficiency virus-related immunosuppression in female sex workers in Abidjan, Ivory Coast.** *J Infect Dis* 1995, **172**(5):1371–1374.
  38. Meda N, Sangaré L, Lankoandé S, Sanou P, Campaore PT, Catraye J, Cartoux M, Soudré RB: **Pattern of sexually transmitted diseases among pregnant women in Burkina Faso, West Africa: potential for a clinical management based on simple approaches.** *Genitourin Med* 1997, **73**:188–193.
  39. Okesola A, Fawole O: **Prevalence of human papillomavirus genital infections in sexually transmitted diseases clinic attendees in Ibadan.** *West Afr J Med* 2000, **19**(3):195–199.
  40. Bakare RA, Oni AA, Umar US, Adewole IF, Shokunbi WA, Fayemiwo SA, Fasina NA: **Pattern of sexually transmitted diseases among commercial sex workers (CSWs) in Ibadan, Nigeria.** *Afr J Med Med Sci* 2002, **31**(3):243–247.
  41. Domfeh AB, Wiredu EK, Adjei AA, Ayeh-Kumi PFK, Adiku TK, Tet-ty Y, Gyasi RK, Armah HB: **Cervical human papillomavirus infection in Accra, Ghana.** *Ghana Med J* 2008, **42**(2):71–78.
  42. Sagay AS, Imade GE, Onwuliri V, Egah DZ, Grigg MJ, Musa J, Thacher TD, Adisa JO, Potts M, Short RV: **Genital tract abnormalities among female sex workers who douche with lemon/lime juice in Nigeria.** *Afr J Reprod Health* 2009, **13**(1):37–45.
  43. Jombo E, Okwori E, Otor G, Odengle E: **Patterns of genital ulcer diseases among HIV/AIDS patients in Benue State, North Central Nigeria.** *Internet J Epidemiol* 2009, **7**:2.
  44. Low A, Clayton T, Konate I, Nagot N, Ouedraogo A, Huet C, Didelot-Rousseau M-N, Michel S, Van de Perre P, Mayaud P, the Yérélon Cohort Study Group: **Genital warts and infection with human immunodeficiency virus in high-risk women in Burkina Faso: a longitudinal study.** *BMC Infect Dis* 2011, **11**:20.
  45. Smith J, Moses S, Hudgens MG, Parker CB, Agot K, Maclean I, Ndinya-Acholi JO, Snijders P, Meijer JF, C. J. L. M, Bailey RC: **Increased risk of HIV acquisition among Kenyan men with human papillomavirus infection.** *JID* 2010, **201**(11):1677–1685.
  46. Tobian AA, Grabowski MK, Kigozi G, Gravitt PE, Eaton KP, Serwadda D, Nalugoda F, Wawer MJ, Quinn TC, Gray RH: **High-risk human papillomavirus prevalence is associated with HIV infection among heterosexual men in Rakai, Uganda.** *Sex Transm Infect* 2012, **89**(2):122–127.
  47. Machekano R, Bassett M, Zhou P, Mbizvo M, Latif A, Katzenstein D: **Report of sexually transmitted diseases by HIV infected man during follow up: time to target the HIV infected?** *Sex Transm Inf* 2000, **76**:186–192.
  48. Müller EE, Chirwa TF, Lewis DA: **Human papillomavirus infection in heterosexual South African men attending sexual health services: associations between HPV and HIV serostatus.** *Sex Transm Infect* 2010, **86**:175–180.
  49. Wade AS, Kane CT, Diallo PA, Diop AK, Gueye K, Mboup S, Ndoeye I, Lagarde E: **HIV infection and sexually transmitted infections among men who have sex with men in Senegal.** *AIDS* 2005, **19**:2133–2140.
  50. Lavreys L, Rakwar J, Thompson M, Jackson D, Mandaliya K, Chohan BH, Bwayo JJ, Ndinya-Ochola JO, Kreiss JK: **Effect of circumcision on incidence of HIV-1 and other sexually transmitted diseases: A prospective cohort study of trucking company employees in Kenya.** *J Infect Dis* 1999, **180**(2):330–336.
  51. Ozumba B, Megafu U: **Pattern of vulval warts at the University of Nigeria teaching hospital, Enugu, Nigeria.** *Int J Gynecol Obstet* 1991, **34**:347–352.
  52. Wright C, Taylor L, Cooper K: **HPV typing of vulvovaginal condylomata in children.** *S Afr Med J* 1995, **85**(10 Suppl):1096–1101.
  53. Firnhaber C, Sello M, Maskew M, Williams S, Schulze D, Williamson AL, Allan B, Lewis D: **Human papillomavirus types in HIV seropositive men with penile warts in Johannesburg, South Africa.** *Int J STD AIDS* 2011, **22**(2):107–109.
  54. Patel H, Wagner M, Singhal P, Kothari S: **Systematic review of the incidence and prevalence of genital warts.** *BMC Infect Dis* 2013, **13**:39.
  55. Silverberg M, Ahdieh LM, Anastos K, Burk R, Cu-Uvin SD, Greenblatt RM, Klein RS, Massad S, Minkoff H, Murderspach L, Palefsky J, Piessens E, Schuman P, Watts H, Shah KV: **The impact of HIV infection and immunodeficiency on human papillomavirus type 6 or 11 infection and on genital warts.** *Sex Transm Dis* 2002, **29**:427–435.
  56. Strickler H, Burk RD, Fazzari M: **Natural history and possible reactivation of human papillomavirus type 6 or 11 infection and on genital warts.** *Sex Transm Dis* 2002, **29**:427–435.
  57. Massad L, Silverberg M, Springer G, Minkoff H, Hessel N, Palefsky JM, Strickler HD, Levine AM, Sacks HS, Moxley M, Heather WD: **Effect of antiretroviral therapy on incidence of genital warts and vulvar neoplasia among women with human immunodeficiency virus.** *Am J Obstet Gynecol* 2004, **190**:1241–1248.
  58. Nasti G, Martellota F, Berretta M, Mena M, Fasan M, Di Perri G: **Impact of highly active anti retroviral therapy on the presenting features and outcome of patients with acquired immunodeficiency syndrome-related Kaposi sarcoma.** *Cancer* 2003, **11**:2440–2446.
  59. Heard I, Tassie J, Schmitz V, Mandelbrot L, Kazatchkine M, Orth G: **Increased risk of cervical disease among human immunodeficiency virus-infected women with severe immunosuppression and high human papillomavirus load(1).** *Obstet Gynecol* 2000, **96**:403–409.
  60. Manhart L, Koutsky L: **Do condoms prevent genital HPV infection, external genital warts or cervical neoplasia?** *Sex Transm Dis* 2002, **29**(11):725–735.
  61. Aubin F, Prétet JL, Jacquard AC, Saunier M, Carcopino X, Jaroud F, Pradat P, Soubeyrand B, Leocmach Y, Mouglin C, Riethmuller D, EDITH Study Group: **Human papillomavirus genotype distribution in external acuminata condylomata: a Large French National Study (EDITH IV).** *Clin Infect Dis* 2008, **47**:610–615.
  62. Vandepapeliere P, Barraso R, Meijer CJ, Walboomers JM, Wettendorff M, Stanberry LR, Lacey CJ: **Randomized controlled trial of an adjuvanted human papillomavirus type 6 L2E7 vaccine: infection of external genital warts with multiple HPV types and failure of therapeutic vaccination.** *J Infect Dis* 2005, **192**:2099–2107.
  63. Wen L, Estcourt C, Simpson J, Mindel A: **Risk factors for acquisition of genital warts: are condoms protective?** *Sex Transm Infect* 1999, **75**:312–316.
  64. Moscicki A, Hills N, Shiboski S, Powell K, Jay N, Hanson E, Miller S, Clayton L, Farhat S, Broering J, Darragh T, Palefsky J: **Risks for incident human papillomavirus infection and low-grade squamous intraepithelial lesion development in young females.** *JAMA* 2001, **285**:2995–3002.
  65. Watts DH, Fazzari M, Minkoff H, Hillier SL, Sha B, Glesby M, Levine AM, Burk R, Palefsky JM, Moxley M, Ahdieh-Grant L, Strickler HD: **Effects of bacterial vaginosis and other genital infections on the natural history of human papillomavirus infection in HIV-1 infected and high-risk HIV-1 uninfected women.** *J Infect Dis* 2005, **191**(7):1129–1139.
  66. Persson G, Dahlöf LG, Krantz I: **Physical and psychological effects of anogenital warts in female patients.** *Sex Transm Dis* 1993, **20**:10–13.

doi:10.1186/1750-9378-8-27

**Cite this article as:** Banura et al.: Prevalence, incidence and risk factors for anogenital warts in Sub Saharan Africa: a systematic review and meta analysis. *Infectious Agents and Cancer* 2013 **8**:27.